

THE SPECTRUM OF THE LAPLACE-BELTRAMI OPERATOR ON THE TWO-DIMENSIONAL TORUS IN ADIABATIC LIMIT

ANDREY A. YAKOVLEV

ABSTRACT. We obtain an asymptotic formula for the eigenvalue distribution function of the Laplace-Beltrami operator on the two-dimensional torus in the adiabatic limit given by a Kronecker foliation. Related problems in number theory are discussed.

INTRODUCTION

Consider the two-dimensional torus $\mathbb{T}^2 = \mathbb{R}^2/\mathbb{Z}^2$ with the coordinates $(x, y) \in \mathbb{R}^2$, considered modulo integer translations, and the Euclidean metric g on $\mathbb{T}^2$:

$$g = dx^2 + dy^2.$$

Let $\tilde{X}$ be the vector field on $\mathbb{R}^2$ given by

$$\tilde{X} = \frac{\partial}{\partial x} + \alpha \frac{\partial}{\partial y},$$

where $\alpha \in \mathbb{R}$. Since $\tilde{X}$ is translation invariant, it determines a vector field X on $\mathbb{T}^2$. The vector field X defines a foliation $\mathcal{F}$ on $\mathbb{T}^2$. The leaves of $\mathcal{F}$ are the images of the parallel lines $\tilde{L} = \{(x_0 + t, y_0 + t\alpha) : t \in \mathbb{R}, (x_0, y_0) \in \mathbb{R}^2\}$ with the slope α under the projection $\mathbb{R}^2 \rightarrow \mathbb{T}^2$.

In the case when α is rational, all leaves of $\mathcal{F}$ are closed and are circles, and $\mathcal{F}$ is given by the fibers of a fibration of $\mathbb{T}^2$ over $\mathbb{S}^1$. In the case when α is irrational, all leaves of $\mathcal{F}$ are everywhere dense in $\mathbb{T}^2$.

Let $F = T\mathcal{F}$ be the tangent bundle to $\mathcal{F}$:

$$F_{(x,y)} = \{(t, \alpha t) : t \in \mathbb{R}\} = \langle (1, \alpha) \rangle \subset T_{(x,y)}\mathbb{T}^2,$$

and $H = F^\perp = \langle (-\alpha, 1) \rangle$ is the orthogonal complement for F . Thus, we have the decomposition of the bundle $T\mathbb{T}^2$ into direct sum $T\mathbb{T}^2 = F \oplus H$

$$T_{x,y}\mathbb{T}^2 \ni (X, Y) = \frac{X + \alpha Y}{1 + \alpha^2}(1, \alpha) + \frac{-\alpha X + Y}{1 + \alpha^2}(-\alpha, 1)$$

Supported by Russian Foundation of Basic Research (grant no. 06-01-00208).

and the corresponding decomposition of the Riemannian metric $g = g_F + g_H$. Define a one-parameter family g_h of Riemannian metrics on $\mathbb{T}^2$ by the formula

$$g_h = g_F + h^{-2}g_H, \quad 0 < h \leq 1.$$

In the coordinates (x, y) , the Riemannian metric g_h is given by

$$g_h = \frac{1 + h^{-2}\alpha^2}{1 + \alpha^2}dx^2 + 2\alpha\frac{1 - h^{-2}}{1 + \alpha^2}dxdy + \frac{\alpha + h^{-2}}{1 + \alpha^2}dy^2.$$

Recall that, for an arbitrary n -dimensional manifold M with a Riemannian metric g , an elliptic, second order, differential operator $\Delta = \Delta_g$ in the space $C_0^\infty(M)$, which is called the Laplace-Beltrami operator, is defined. In local coordinates, this operator is given by the formula

$$\Delta = -\frac{1}{\sqrt{g}} \sum_{i,j=1}^n \frac{\partial}{\partial x^i} \left(g^{ij} \sqrt{g} \frac{\partial}{\partial x^j} \right),$$

where (g_{ij}) is the matrix of the Riemannian metric, g is the determinant of the matrix (g_{ij}) , (g^{ij}) is the inverse of the matrix (g_{ij}) .

In our case, the Laplace operator, defined by the metric g_h , has the form

$$\Delta_h = -\frac{1}{1 + \alpha^2} \left(\frac{\partial}{\partial x} + \alpha \frac{\partial}{\partial y} \right)^2 - \frac{h^2}{1 + \alpha^2} \left(-\alpha \frac{\partial}{\partial x} + \frac{\partial}{\partial y} \right)^2 = \Delta_F + h^2 \Delta_H,$$

where Δ_F is the tangential Laplace operator:

$$\Delta_F = -\frac{1}{1 + \alpha^2} \left(\frac{\partial}{\partial x} + \alpha \frac{\partial}{\partial y} \right)^2,$$

and Δ_H is the transverse Laplace operator:

$$\Delta_H = -\frac{h^2}{1 + \alpha^2} \left(-\alpha \frac{\partial}{\partial x} + \frac{\partial}{\partial y} \right)^2.$$

The main result of the paper is the calculation of the asymptotics of the spectrum distribution function $N_h(\lambda)$ of the operator Δ_h for a fixed $\lambda \in \mathbb{R}$ and for $h \rightarrow 0$.

Theorem 1. *The following asymptotic formula for the spectrum distribution function $N_h(\lambda)$ of the operator Δ_h for a fixed $\lambda \in \mathbb{R}$ holds:*

1. For $\alpha \notin \mathbb{Q}$,

$$N_h(\lambda) = \frac{1}{4\pi} h^{-1} \lambda + o(h^{-1}), \quad h \rightarrow 0.$$

2. For $\alpha \in \mathbb{Q}$ of the form $\alpha = \frac{p}{q}$, where $\text{G.C.D.}(p, q) = 1$,

$$N_h(\lambda) = h^{-1} \sum_{k: \frac{4\pi^2}{p^2+q^2} k^2 < \lambda} \frac{1}{\pi \sqrt{p^2+q^2}} (\lambda - \frac{4\pi^2}{p^2+q^2} k^2)^{1/2} + o(h^{-1}), \quad h \rightarrow 0.$$

We will give two proofs of Theorem 1. The first proof is based on a general result on the asymptotic behavior of the spectrum distribution function of the Laplace operator on a compact manifold equipped with a Riemannian foliation in the adiabatic limit, obtained in the work [2]. The second proof is considerably simpler and uses only elementary facts of analysis.

Regarding various notions and facts, connected with the foliation theory see, for instance, the survey paper [3] and the bibliography cited there.

The results of the paper are announced in [4].

The author is grateful to Kordyukov Yu.A. for the setting of the problem and useful discussions.

1. ADIABATIC LIMITS FOR RIEMANNIAN FOLIATIONS

Let $(M, \mathcal{F})$ be a closed foliated manifold, $\dim M = n$, $\dim \mathcal{F} = p$, $p + q = n$, equipped with a Riemannian metric g_M . Let $F = T\mathcal{F}$ be the integrable distribution of tangent p -planes to $\mathcal{F}$ in TM , and $H = F^\perp$ the orthogonal complement to F . Thus, we have a direct sum decomposition of TM :

$$TM = F \oplus H.$$

This decomposition induces a decomposition of the metric $g_M = g_F + g_H$. Define a one-parameter family g_h of metrics on M by the formula

$$g_h = g_F + h^{-2} g_H, \quad 0 < h \leq 1.$$

For any $h > 0$ we consider the Laplace-Beltrami operators on functions, determined by the metric g_h :

$$\Delta_h = d_{g_h}^* d,$$

where $d : C^\infty(M) \rightarrow C^\infty(M, \Lambda^1 T^*M)$ is the de Rham differential, $d_{g_h}^*$ is the adjoint operator of d with respect to the inner products, defined by the metric g_h . The operator Δ_h is a self-adjoint, elliptic, differential operator with the positive definite, scalar principal symbol in the Hilbert space $L^2(M, g_h)$ of square integrable differential forms on M , endowed with the inner product, induced by the metric g_h . According to the standard perturbation theory there exists a countable family of analytic functions $\lambda_i(h)$ such that, for any $h > 0$,

$$\text{spec } \Delta_h = \{\lambda_i(h) : i = 0, 1, \dots\},$$

taking into account multiplicities. Introduce the eigenvalue distribution function of the operator Δ_h :

$$N_h(\lambda) = \#\{i : \lambda_i(h) \leq \lambda\}$$

Suppose that the foliation $\mathcal{F}$ is Riemannian and g_M is a bundle-like Riemannian metric. Let dx be the Riemannian volume form on M .

Denote by Δ_F the tangential Laplace operator on $C^\infty(M)$ and by Δ_L its restriction to the leaf L of $\mathcal{F}$. The operator Δ_L coincides with the Laplace operator on L , defined by the induced metric of the leaf.

Let G be the holonomy groupoid of $\mathcal{F}$. Let λ_L denote the Riemannian volume form on a leaf L given by the induced metric, and λ^x denote its lift to the holonomy covering G^x for any $x \in M$.

Denote by $e_\lambda(\gamma), \gamma \in G, \lambda \in \mathbb{R}$ the tangential kernel of the spectral projection of the tangentially elliptic operator Δ_F , corresponding to the semi-axis $(-\infty, \lambda)$. For every $x \in M$ the function $e_\lambda(\gamma^{-1}\gamma'), (\gamma, \gamma') \in G^x$ coincides with the kernel of the spectral projection of the operator Δ_{G^x} , which is the lift of Δ_L to the holonomy covering G^x , relative to the Riemannian volume form λ^x . Let $e_\lambda(x), x \in M, \lambda \in \mathbb{R}$ be the restriction of $e_\lambda(\gamma)$ to M . For any $\lambda \in \mathbb{R}$, the function e_λ is a bounded measurable function on M .

The spectrum distribution function $N_{\mathcal{F}}(\lambda)$ of Δ_F is defined by the formula

$$N_{\mathcal{F}}(\lambda) = \int_M e_\lambda(x) dx.$$

Theorem 2. *Let $(M, \mathcal{F})$ be a Riemannian foliation, equipped with a bundle-like metric g_M . The following asymptotic formula holds:*

(1)

$$N_h(\lambda) = h^{-q} \frac{(4\pi)^{-q/2}}{\Gamma((q/2) + 1)} \int_{-\infty}^{\lambda} (\lambda - \tau)^{q/2} d_\tau N_{\mathcal{F}}(\tau) + o(h^{-q}), \quad h \rightarrow 0.$$

2. PROOF OF THEOREM 1

2.1. The first proof. In this section we derive the asymptotic formula for $N_h(\lambda)$ on the torus from Theorem 2.

The induced metric on the leaf $x = x_0 + t, y = y_0 + \alpha t$ through a point (x_0, y_0) has the form $g_{\mathcal{F}} = (1 + \alpha^2)dt^2$. Thus, the leafwise Laplace operator is given by

$$\Delta_F = -\frac{1}{1 + \alpha^2} \left(\frac{\partial}{\partial x} + \alpha \frac{\partial}{\partial y} \right)^2.$$

1. $\alpha \notin \mathbb{Q}$. In this case $G = \mathbb{T}^2 \times \mathbb{R}$. The source and the range maps $s, r : G \rightarrow \mathbb{T}^2$ are defined for any $\gamma = (x, y, t) \in G$ by $s(\gamma) =$

$(x - t, y - \alpha t)$ and $r(\gamma) = (x, y)$. For any $(x, y) \in \mathbb{T}^2$, the set $G^{(x,y)}$ coincides with the leaf $L_{(x,y)}$ through (x, y) and is diffeomorphic to $\mathbb{R}$:

$$L_{(x,y)} = \{(x - t, y - \alpha t) : t \in \mathbb{R}\}.$$

The Riemannian volume form $\lambda^{(x,y)}$ on $L_{(x,y)}$ equals $\sqrt{1 + \alpha^2} dt$. Finally, the restriction of the operator Δ_F to each leaf $L_{(x,y)}$ coincides with the operator

$$A = -\frac{1}{1 + \alpha^2} \frac{d^2}{dt^2},$$

acting in the space $L^2(\mathbb{R}, \sqrt{1 + \alpha^2} dt)$.

Let us find the spectral projections of A in $L^2(\mathbb{R}, \sqrt{1 + \alpha^2} dt)$. The following equality is valid:

$$A_1 U(\xi) = F A F^{-1} U(\xi) = \frac{|\xi|^2}{1 + \alpha^2} U(\xi), \quad U \in C_0^\infty(\mathbb{R}),$$

where F is the Fourier transform. A similar equality holds for the spectral projections:

$$\chi_\lambda(A_1) = F \chi_\lambda(A) F^{-1},$$

where χ_λ is the characteristic function of the semi-axis $(-\infty, \lambda)$. It is easy to see that the operator $\chi_\lambda(A_1)$ is the multiplication operator by the the function

$$\chi_\lambda \left(\frac{|\xi|^2}{1 + \alpha^2} \right) = \begin{cases} 1, & \text{if } \frac{|\xi|^2}{1 + \alpha^2} < \lambda, \\ 0, & \text{if } \frac{|\xi|^2}{1 + \alpha^2} \geq \lambda. \end{cases}$$

Thus, we get that the spectral projection $\chi_\lambda(A)$ of the operator A is of the form:

$$\begin{aligned} \chi_\lambda(A) U(t) &= F^{-1} \chi_\lambda \left(\frac{|\xi|^2}{1 + \alpha^2} \right) F U(t) \\ &= \frac{1}{2\pi} \int_{\mathbb{R}^2} \exp[i(t - t_1)\xi] \chi_\lambda \left(\frac{|\xi|^2}{1 + \alpha^2} \right) U(t) dt_1 d\xi. \end{aligned}$$

The kernel of the spectral projection $\chi_\lambda(A)$ of A in the space $L^2(\mathbb{R})$ (relative to the Riemannian volume form $\sqrt{1 + \alpha^2} dt$) is given by the formula

$$E_\lambda(t, t_1) = \frac{1}{2\pi\sqrt{1 + \alpha^2}} \int_{\mathbb{R}} \exp[i(t - t_1)\xi] \chi_\lambda \left(\frac{|\xi|^2}{1 + \alpha^2} \right) d\xi.$$

The tangential kernel e_λ of the operator Δ_F is related with E_λ as follows. For any $\gamma = (x, y, t) \in G = \mathbb{T}^2 \times \mathbb{R}$,

$$e_\lambda(\gamma) = E_\lambda(0, t) = \frac{1}{2\pi\sqrt{1 + \alpha^2}} \int_{\mathbb{R}} \exp(-it\xi) \chi_\lambda \left(\frac{|\xi|^2}{1 + \alpha^2} \right) d\xi.$$

The restriction of the tangential kernel e_λ to $\mathbb{T}^2$ is given by

$$e_\lambda(x, y) = E_\lambda(0, 0) = \frac{1}{2\pi\sqrt{1+\alpha^2}} \int_{\mathbb{R}} \chi_\lambda \left(\frac{|\xi|^2}{1+\alpha^2} \right) d\xi = \frac{1}{\pi} \sqrt{\lambda}, \quad \lambda > 0.$$

We get that the spectrum distribution function $N_{\mathcal{F}}(\lambda)$ of operator Δ_F has the form:

$$N_{\mathcal{F}}(\lambda) = \int_{\mathbb{T}^2} e_\lambda(x, y) dx dy = \frac{1}{\pi} \sqrt{\lambda}, \quad \lambda > 0.$$

By Theorem 2, we obtain

$$N_h(\lambda) = h^{-1} \frac{1}{\pi} \int_{-\infty}^{\lambda} (\lambda - \tau)^{1/2} d_\tau N_{\mathcal{F}}(\tau) + o(h^{-1}), \quad h \rightarrow 0.$$

Hence,

$$N_h(\lambda) = \frac{1}{4\pi} h^{-1} \lambda + o(h^{-1}), \quad h \rightarrow 0.$$

2. $\alpha \in \mathbb{Q}$. Let us consider $\alpha = \frac{p}{q}$, where p and q are coprime. In this case, the holonomy groupoid is $\mathbb{T}^2 \times (\mathbb{R}/q\mathbb{Z})$. The leaf through an arbitrary point (x_0, y_0) is the circle $\{(x_0 + t, y_0 + \alpha t) : t \in \mathbb{R}/q\mathbb{Z}\}$ of length $l = \sqrt{p^2 + q^2}$.

The restriction of the operator Δ_F to each $L_{(x,y)}$ coincides with the operator

$$A = -\frac{1}{1+\alpha^2} \frac{d^2}{dt^2},$$

acting in the space $L^2(\mathbb{R}/q\mathbb{Z}, \sqrt{1+\alpha^2} dt)$.

This operator has a complete orthogonal system of eigenfunctions $U_j(t)$ the corresponding eigenvalues λ_j :

$$\lambda_j = \frac{1}{1+\alpha^2} \frac{4\pi^2}{q^2} j^2 = \frac{4\pi^2}{p^2 + q^2} j^2, \quad U_j(t) = \exp\left[\frac{2\pi i}{q} jt\right], \quad j \in \mathbb{Z}.$$

Any function $U(t) \in L^2(\mathbb{R}/q\mathbb{Z}, \sqrt{1+\alpha^2} dt)$ can be represented as the sum of a series:

$$U(t) = \frac{1}{q} \sum_{k=-\infty}^{\infty} \left(\int_0^q U(t_1) \exp\left[-\frac{2\pi i}{q} kt_1\right] dt_1 \right) \exp\left[\frac{2\pi i}{q} kt\right]$$

Therefore, the spectral projection $\chi_\lambda(A)$ of A is given by the formula:

$$\chi_\lambda(A)U(t) = \frac{1}{q} \sum_{k: \frac{4\pi^2}{p^2+q^2} k^2 < \lambda} \left(\int_0^q U(t_1) \exp\left[-\frac{2\pi i}{q} kt_1\right] dt_1 \right) \exp\left[\frac{2\pi i}{q} kt\right]$$

The kernel of the projection $\chi_\lambda(A)$ in $L^2(\mathbb{R}/q\mathbb{Z}, \sqrt{1+\alpha^2} dt)$ is given by the formula

$$E_\lambda(t, t_1) = \frac{1}{\sqrt{p^2 + q^2}} \sum_{k: \frac{4\pi^2}{p^2 + q^2} k^2 < \lambda} \exp\left[\frac{2\pi i}{q} k(t - t_1)\right].$$

The tangential kernel e_λ of Δ_F is related with E_λ in the following way. For any $\gamma = (x, y, t) \in G = \mathbb{T}^2 \times (\mathbb{R}/q\mathbb{Z})$,

$$e_\lambda(\gamma) = E_\lambda(0, t) = \frac{1}{\sqrt{p^2 + q^2}} \sum_{k: \frac{4\pi^2}{p^2 + q^2} k^2 < \lambda} \exp\left[-\frac{2\pi i}{q} kt\right].$$

The restriction e_λ to $\mathbb{T}^2$ has the form:

$$e_\lambda(x, y) = E_\lambda(0, 0) = \frac{1}{\sqrt{p^2 + q^2}} \#\{k : \frac{4\pi^2}{p^2 + q^2} k^2 < \lambda\}.$$

We get that, in the case of a rational α , the spectrum distribution function $N_{\mathcal{F}}(\lambda)$ of Δ_F is of the form:

$$N_{\mathcal{F}}(\lambda) = \int_{\mathbb{T}^2} e_\lambda(x, y) dx dy = \frac{1}{\sqrt{p^2 + q^2}} \#\{k : \frac{4\pi^2}{p^2 + q^2} k^2 < \lambda\}.$$

By Theorem 2, we obtain for $h \rightarrow 0$

$$\begin{aligned} N_h(\lambda) &= h^{-1} \frac{1}{\pi} \int_{-\infty}^{\lambda} (\lambda - \tau)^{1/2} d_\tau N_{\mathcal{F}}(\tau) + o(h^{-1}) \\ &= h^{-1} \frac{1}{\pi \sqrt{p^2 + q^2}} \sum_{k: \frac{4\pi^2}{p^2 + q^2} k^2 < \lambda} \left(\lambda - \frac{4\pi^2}{p^2 + q^2} k^2\right)^{1/2} + o(h^{-1}). \end{aligned}$$

2.2. The second proof. The operator Δ_h has a complete orthogonal system of eigenfunctions

$$u_{kl}(x, y) = e^{2\pi i(kx + ly)}, \quad (x, y) \in \mathbb{T}^2,$$

with the corresponding eigenvalues

$$\lambda_{kl} = (2\pi)^2 \left(\frac{1}{1 + \alpha^2} (k + \alpha l)^2 + \frac{h^2}{1 + \alpha^2} (-\alpha k + l)^2 \right), \quad k, l \in \mathbb{Z}.$$

The eigenvalue distribution function of Δ_h has the form

$$\begin{aligned} N_h &= \#\{(k, l) \in \mathbb{Z}^2 : \\ &\quad (2\pi)^2 \left(\frac{1}{1 + \alpha^2} (k + \alpha l)^2 + \frac{h^2}{1 + \alpha^2} (-\alpha k + l)^2 \right) < \lambda\}. \end{aligned}$$

Thus we come to the following problem of number theory:

Problem 1. Find the asymptotics for $h \rightarrow 0$ of the number of integer points in the ellipse

$$\{(\xi, \eta) \in \mathbb{R}^2 : (2\pi)^2 \left(\frac{1}{1+\alpha^2}(\xi + \alpha\eta)^2 + \frac{h^2}{1+\alpha^2}(-\alpha\xi + \eta)^2 \right) < \lambda\}.$$

Let us introduce new coordinates

$$\begin{cases} \xi' = \frac{1}{\sqrt{1+\alpha^2}}\xi + \frac{\alpha}{\sqrt{1+\alpha^2}}\eta, \\ \eta' = -\frac{\alpha}{\sqrt{1+\alpha^2}}\xi + \frac{1}{\sqrt{1+\alpha^2}}\eta. \end{cases}$$

In the new coordinates we will consider a point (ξ', η') to be integer, if it has the form

$$\begin{cases} \xi' = \frac{1}{\sqrt{1+\alpha^2}}n + \frac{\alpha}{\sqrt{1+\alpha^2}}m, \\ \eta' = -\frac{\alpha}{\sqrt{1+\alpha^2}}n + \frac{1}{\sqrt{1+\alpha^2}}m, \end{cases}$$

where $(n, m) \in \mathbb{Z}^2$. Hence, the number of such integer points is determined by the number of all pairs $(n, m) \in \mathbb{Z}^2$, satisfying the following conditions:

$$\begin{cases} \left| \frac{1}{\sqrt{1+\alpha^2}}n + \frac{\alpha}{\sqrt{1+\alpha^2}}m \right| < \frac{\sqrt{\lambda}}{2\pi}, \\ \left| -\frac{\alpha}{\sqrt{1+\alpha^2}}n + \frac{1}{\sqrt{1+\alpha^2}}m \right| < h^{-1} \sqrt{\frac{\lambda}{(2\pi)^2} - \left(\frac{1}{\sqrt{1+\alpha^2}}n + \frac{\alpha}{\sqrt{1+\alpha^2}}m \right)^2}. \end{cases}$$

It is clear that, as $h \rightarrow 0$, for $\alpha \in \mathbb{R}/\mathbb{Q}$ the projections of integer points to the ξ' -axes form an everywhere dense set, but for $\alpha \in \mathbb{Q}$ a discrete one.

Suppose that $\alpha \in \mathbb{Q}$. Without loss of generality, we will assume $\alpha = \frac{p}{q}$, where $\text{G.C.D.}(p, q) = 1$. By straightforward computations, it is easy to show that a point is integer, if has the form:

$$\left(\frac{k}{\sqrt{p^2 + q^2}}, \frac{-pn_0 + qm_0 - (p^2 + q^2)l}{\sqrt{p^2 + q^2}} \right),$$

where (n_0, m_0) is an integer point determined by $k, l \in \mathbb{Z}$ is responsible for the number of integer point over ξ' . Since we are looking for the number of integer points in the ellipse, we get the following conditions on l :

$$\begin{aligned} -h^{-1} \sqrt{\frac{\lambda}{4\pi^2} - \frac{k^2}{p^2 + q^2}} + \frac{-pn_0 + qm_0}{\sqrt{p^2 + q^2}} &< \sqrt{p^2 + q^2}l \\ &< h^{-1} \sqrt{\frac{\lambda}{4\pi^2} - \frac{k^2}{p^2 + q^2}} + \frac{-pn_0 + qm_0}{\sqrt{p^2 + q^2}}. \end{aligned}$$

Thus, the number of integer points over ξ' is equal to

$$2 \frac{h^{-1}}{\sqrt{p^2 + q^2}} \sqrt{\frac{\lambda}{4\pi^2} - \frac{k^2}{p^2 + q^2}}.$$

We obtain that in the case of rational $\alpha = p/q$ the asymptotics of the eigenvalue distribution function $N_h(\lambda)$ as $h \rightarrow 0$ is given by

$$N_h(\lambda) = h^{-1} \sum_{k: \frac{4\pi^2}{p^2+q^2} k^2 < \lambda} \frac{1}{\pi \sqrt{p^2 + q^2}} (\lambda - \frac{4\pi^2}{p^2 + q^2} k^2)^{1/2} + o(h^{-1}).$$

Suppose that $\alpha \notin \mathbb{Q}$. Let us the heat operator $e^{-t\Delta_h}$ defined by the Laplace operator Δ_h in $\mathbb{R}^2$. For any $u \in L^2(\mathbb{R}^2)$ the function $u(t) = e^{-t\Delta_h} u$ is a solution of the heat equation

$$\frac{\partial u}{\partial t} = -\Delta_h u, \quad t > 0,$$

with initial conditions $u(0) = u$. It can be shown that the kernel of $e^{-t\Delta_h}$ on $\mathbb{R}^2$ has the form

$$\begin{aligned} H_t(x, y, x_1, y_1) &= \\ &= \frac{h^{-1}}{4\pi t} \exp \left[-\frac{(x - x_1 + \alpha(y - y_1))^2 + h^{-2}(-\alpha(x - x_1) + y - y_1)^2}{4t(1 + \alpha^2)} \right]. \end{aligned}$$

Therefore, the kernel of $e^{-t\Delta_h}$ in $\mathbb{T}^2$ is given by

$$\begin{aligned} H_t(x, y, x_1, y_1) &= \frac{h^{-1}}{4\pi t} \sum_{k, l=-\infty}^{+\infty} \exp \left[-\frac{(x - x_1 + k + \alpha(y - y_1 + l))^2}{4t(1 + \alpha^2)} - \right. \\ &\quad \left. - \frac{(-\alpha(x - x_1 + k) + y - y_1 + l)^2}{4th^2(1 + \alpha^2)} \right]. \end{aligned}$$

Accordingly, the trace of $tr(e^{-t\Delta_h})$ in $L^2(\mathbb{T}^2)$ is given by

$$tr(e^{-t\Delta_h}) = \frac{h^{-1}}{4\pi t} \sum_{k, l=-\infty}^{+\infty} e^{-\frac{(k+\alpha l)^2}{4t(1+\alpha^2)} - \frac{(-\alpha k+l)^2}{4th^2(1+\alpha^2)}}.$$

Since the series converges uniformly on $h \in (0, 1]$ and, for any $(k, l) \neq (0, 0)$

$$e^{-\frac{(k+\alpha l)^2}{4t(1+\alpha^2)} - \frac{(-\alpha k+l)^2}{4th^2(1+\alpha^2)}} \longrightarrow 0, \quad h \rightarrow +0,$$

we get

$$\text{tr } e^{-t\Delta_h} \longrightarrow \frac{h^{-1}}{4\pi t}, \quad h \rightarrow +0.$$

From the other side, we have

$$\mathrm{tr} e^{-t\Delta_h} = \int_{-\infty}^{+\infty} e^{-\lambda t} d_\lambda N_h(\lambda),$$

therefore, the proof of Theorem 1 is completed by the next lemma ([5, Lemma 5.2], cf. also [1]).

We say that a function F on the real line is a distribution function if F is a non-decreasing left-continuous function such that

$$\lim_{\lambda \rightarrow -\infty} F(\lambda) = 0.$$

For any distribution function $F(\lambda)$, we denote by $\tilde{F}(t)$ the Laplace transform of this function.

Lemma 3. *Let F_n be a sequence of distribution functions on the real line such that:*

- (a) $F_n(\lambda) = 0, \lambda \leq a$, with the constant a , not depending on n ;
- (b) $|F_n(\lambda)| \leq Ce^{-\varepsilon\lambda}, \lambda \in \mathbb{R}$, with the constants C and ε , not depending on n ;
- (c) there exists $\lim_{n \rightarrow \infty} \tilde{F}_n(t) = \tilde{F}(t)$ for any $t > 0$, where $F(t)$ is a distribution function.

Then $\lim_{n \rightarrow \infty} F_n(\lambda) = F(\lambda)$ at all points of continuity of the function $F(\lambda)$.

REFERENCES

- [1] Avron J., Simon B. Almost periodic Schrödinger operators. II. The integrated density of states; *Duke Math. J.*, **50** (1983), 369–391.
- [2] Kordyukov Yu. A. Adiabatic limits and spectral geometry of foliations. *Math. Ann.* **313** (1999), 763–783.
- [3] Kordyukov Yu. A. Noncommutative geometry of foliations. Preprint math.DG/0504095, 2005.
- [4] Kordyukov Yu. A., Yakovlev A.A. Adiabatic limits and the spectrum of the Laplacian on foliated manifolds, submitted to Proceedings of the Conference “ C^* -algebras and elliptic theory. II” (Bedlewo, Poland, 2006).
- [5] Shubin M.A. The density of states of self-adjoint elliptic operators with almost-periodic coefficients, *Amer. Math. Soc. Trans.* **118** (1982), 307 – 339.

DEPARTMENT OF MATHEMATICS,, UFA STATE AVIATION TECHNICAL UNIVERSITY, 12 K.MARX STR., UFA, 450000, RUSSIA

E-mail address: yakovlevandrey@yandex.ru